

Kybella Post-Care Instructions

1. Do NOT, touch, press, rub, massage, or manipulate the treatment area.
2. Ice the treated areas for the next 24 hours. Place the icepack on the area for 20 minutes and remove the ice pack for 20 minutes. Continue this pattern for 24 hours.
3. Apply (over the counter) Arnica gel to the treatment area to help decrease the bruising, swelling, and discomfort.
4. Take (over the counter) acetaminophen and/or pain reliever if needed to decrease post treatment discomfort.
5. Sleep on your back and with head elevated with 2-3 pillows for the next 3-5 days after treatment.
6. Drink plenty of water and fluids after treatment.
7. Avoid vigorous exercise, sun and heat exposure for 3-5 days after treatment.
8. Avoid Aspirin, Motrin, Ginkgo Biloba, Garlic, Flax Oil, Cod Liver Oil, Vitamin A, Vitamin E or any other essential fatty acids at least 3 days to 1 week after treatment. These items may increase bleeding and bruising.
9. Avoid alcohol, caffeine, Niacin supplement, high-sodium foods, high sugar foods, refined carbohydrates, spicy foods and cigarettes 24-48 hours after your treatment. These items may contribute to increased swelling or irritation.
10. Avoid cosmetic treatments such as injectables, laser, ultrasound, peels, facials or micro-dermabrasion to the Kybella treatment area.
11. It is absolutely normal to feel and see mild to moderate swelling, numbness, hardness to touch, tingling, redness, and tenderness at the site of injections and they can last up to 6-8 weeks. Most swelling will begin to improve after day 5.
12. Please report to The Youth Injector immediately if you develop an asymmetric smile or facial muscle weakness, skin ulceration in the treatment area, difficulty swallowing, or if any existing symptom worsens.

****Don't forget to schedule your next treatment for 4-6 weeks****

